## §558.120

- (ii) *Limitations.* Do not feed to swine weighing more than 75 pounds body weight; do not feed to swine within 10 weeks of slaughter; do not use in complete feeds containing less than 15 percent crude protein.
- (2) Amount per ton. 50 grams (0.0055 percent).
- (i) Indications for use. For control of swine dysentery (vibrionic dysentery, bloody scours, or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by Salmonella choleraesuis); increased rate of weight gain and improved feed efficiency.
- (ii) *Limitations*. Do not feed to swine weighing more than 75 pounds body weight; do not feed to swine within 10 weeks of slaughter; do not use in complete feeds containing less than 15 percent crude protein.

(3) Amount per ton. Carbadox 50 grams (0.0055 percent) plus pyrantel tartrate, 96 grams (0.0106 percent).

- (i) Indications for use. For control of swine dysentery (vibrionic dysentery, bloody scours, or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by Salmonella choleraesuis); aid in the prevention of migration and establishment of large roundworm (Ascaris suum) infections; aid in the prevention of establishment of nodular worm (Oesophagostomum) infections.
- (ii) Limitations. Do not feed to swine over 75 pounds; do not feed within 10 weeks of slaughter; consult a veterinarian before feeding to severely debilitated animals; feed continuously as sole ration. Do not use in complete feeds containing less than 15 percent crude protein.

[40 FR 13959, Mar. 27, 1975, as amended at 40 FR 45164, Oct. 1, 1975; 40 FR 57798, Dec. 12, 1975; 42 FR 761, Jan. 4, 1977; 51 FR 7396, Mar. 3, 1986]

## §558.120 Carbarsone (not U.S.P.).

- (a) *Approvals*. Type A medicated articles: (1) 37.5 percent to 046573 in \$510.600(c) of this chapter.
- (2) 25 percent carbarsone and 5 grams per pound bacitracin (as bacitracin methylene disalicylate) to 046573 in §510.600(c) of this chapter.
- (b) Related tolerances. See §556.60 of this chapter.

- (c) Conditions of use. (1) It is used for turkeys as follows:
- (i) *Grams per ton.* 227 to 340.5 (0.025 to 0.0375 percent).
- (a) Indications for use. As an aid in the prevention of blackhead.
- (b) Limitations. Feed continuously beginning 2 weeks before blackhead is expected and continue as long as prevention is needed; withdraw 5 days before slaughter; as sole source of organic arsenic.
- (ii) *Grams per ton.* 227 to 340.5 (0.025 to 0.0375 percent) carbarsone plus 10 grams per ton bacitracin from bacitracin methylene disalicylate.
- (a) Indications for use. As an aid in the prevention of blackhead; for increased rate of weight gain.
- (b) Limitations. Feed continuously beginning 2 weeks before blackhead is expected and continue as long as prevention is needed; withdraw 5 days before slaughter; as sole source of organic arsenic.
- (iii) *Grams per ton.* 227 to 340.5 (0.025 to 0.0375 percent) carbarsone plus 4 to 45 grams per ton bacitracin from bacitracin zinc.
- (a) Indications for use. As an aid in the prevention of blackhead, increased rate of weight gain, and improved feed efficiency.
- (b) Limitations. Feed continuously as sole ration. Withdraw 5 days before slaughter. As sole source of organic arsenic; as bacitracin zinc provided by No. 010042 in §510.600(c) of this chapter.
- (2) Carbarsone (not U.S.P.) may be used in accordance with the provisions of this section in the combinations provided as follows:
- (i) Zoalene in accordance with §558.680.
  - (ii) Amprolium as in §558.55.
  - (iii) Bambermycins as in §558.95.

[41 FR 10995, Mar. 15, 1976, as amended at 42 FR 18617, Apr. 8, 1977; 46 FR 46797, Sept. 22, 1981; 48 FR 2758, Jan. 21, 1983; 51 FR 7396, Mar. 3, 1986; 52 FR 2687, Jan. 26, 1987; 53 FR 20843, June 7, 1988; 57 FR 7652, Mar. 4, 1992; 61 FR 515, Jan. 8, 1996]

## §558.128 Chlortetracycline.

(a) Approvals. Type A medicated articles: 10 to 100 grams per pound chlortetracycline to 010042 in §510.600(c) of this chapter; 35 grams chlortetracycline with 7.7 percent (35